

vec, inversevec, diag, svd from linearalgebra

ElementSets from MeshConnectivity

NeighborVerticesInFace, Faces, Vertices, VertexOneRing, OrientedVertices from TetrahedronNeighborhoods(M)

$M : \text{CellMesh}$
 $\bar{x}_i \in \mathbb{R}^3$ rest pos
 $x_i \in \mathbb{R}^3$ current pos
 $bx_j \in \mathbb{Z}$, index boundary indices
 $bp_j \in \mathbb{R}^3$ boundary positions
 $w \in \mathbb{R}$ penalty
 $\varepsilon \in \mathbb{R}$ eps
 $psd : \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{p \times p}$, sparse

$V, E, F, C = \text{ElementSets}(M)$

$\text{vol}_{i,j,k,l} = \frac{1}{6} | [\bar{x}_j - \bar{x}_i \quad \bar{x}_k - \bar{x}_i \quad \bar{x}_l - \bar{x}_i] |$ where $i, j, k, l \in V$
 $m_r(s) = [\bar{x}_b - \bar{x}_a \quad \bar{x}_c - \bar{x}_a \quad \bar{x}_d - \bar{x}_a]$

where

$s \in C$

$a, b, c, d = \text{OrientedVertices}(s)$

$S(s, x) = \begin{cases} \infty & \text{if } |m| \leq 0 \\ \text{vol}_{a,b,c,d} (\|J\|^2 + \|J^{-1}\|^2) & \text{otherwise} \end{cases}$

where

$s \in C$

$x_i \in \mathbb{R}^3$

$a, b, c, d = \text{OrientedVertices}(s)$

$m = [x_b - x_a \quad x_c - x_a \quad x_d - x_a]$

$J = mm_r(s)^{-1}$

$EXPS(s, x) = \begin{cases} \infty & \text{if } |m| \leq 0 \\ \text{vol}_{a,b,c,d} e^{\frac{1}{2}(\|J\|^2 + \|J^{-1}\|^2)} & \text{otherwise} \end{cases}$

where

$s \in C$

$x_i \in \mathbb{R}^3$

$a, b, c, d = \text{OrientedVertices}(s)$

$m = [x_b - x_a \quad x_c - x_a \quad x_d - x_a]$

$J = mm_r(s)^{-1}$

$AMIPS(s, x) = \begin{cases} \infty & \text{if } |m| \leq 0 \\ \text{vol}_{a,b,c,d} e^{\frac{1}{2} \left(\frac{\|J\|^2}{|J|} + \frac{1}{2} (|J| + |J^{-1}|) \right)} & \text{otherwise} \end{cases}$

where

$s \in C$

$x_i \in \mathbb{R}^3$

$a, b, c, d = \text{OrientedVertices}(s)$

$m = [x_b - x_a \quad x_c - x_a \quad x_d - x_a]$

$J = mm_r(s)^{-1}$

$CAMIPS(s, x) = \begin{cases} \infty & \text{if } |m| \leq 0 \\ \text{vol}_{a,b,c,d} \left(\frac{\|J\|^2}{|J|^{\frac{2}{3}}} \right) & \text{otherwise} \end{cases}$

where

$s \in C$

$x_i \in \mathbb{R}^3$

$a, b, c, d = \text{OrientedVertices}(s)$

$m = [x_b - x_a \quad x_c - x_a \quad x_d - x_a]$

$J = mm_r(s)^{-1}$

$E2 = w \sum_j \|bp_j - x_{bx_j}\|^2$

$e = \sum_{i \in C} S_i(x) + E2$

$G = \frac{\partial e}{\partial x}$

$H = \sum_{i \in C} psd \left(\frac{\partial^2 S_i(x)}{\partial x^2} \right) + psd \left(\frac{\partial^2 E2}{\partial x^2} \right)$